



TITLE:

ON RICCATI INEQUALITY(Recent Developments in Theory of Operators and Its Applications)

AUTHOR(S):

Fujii, Jun Ichi; Fujii, Masatoshi; Nakamoto, Ritsuo

CITATION:

Fujii, Jun Ichi ...[et al]. ON RICCATI INEQUALITY(Recent Developments in Theory of Operators and Its Applications). 数理解析研究所講究録 2007, 1535: 31-36

ISSUE DATE:

2007-02

URL:

<http://hdl.handle.net/2433/59000>

RIGHT:

ON RICCATI INEQUALITY (RICCATI 不等式について)

Jun Ichi Fujii (藤井淳一, 大阪教育大学)
Masatoshi Fujii (藤井正俊, 大阪教育大学)
Ritsuo Nakamoto (中本律男, 茨城大学)

1. Introduction.

As stated in our preceding discussion, the algebraic Riccati equation

$$X^*B^{-1}X - T^*X - X^*T = C \quad (B, C; \text{positive definite matrices})$$

has solutions given by $X = W + BT$ for some solution W of

$$W^*B^{-1}W = C + T^*BT$$

because

$$X^*B^{-1}X - T^*X - X^*T = W^*B^{-1}W - T^*BT.$$

Namely the operator equation

$$(1) \quad X^*B^{-1}X = C$$

is essential, so we call it the Riccati equation.

Related to this, we recall Ando's definition of operator geometric mean [2]: For positive operators B, C on a Hilbert space,

$$(2) \quad B \sharp C = \max\{X \geq 0; \begin{pmatrix} B & X \\ X & C \end{pmatrix} \geq 0\}.$$

is called the geometric mean of B and C . If B is invertible, it is expressed by

$$(3) \quad B \sharp C = B^{\frac{1}{2}}(B^{-\frac{1}{2}}CB^{-\frac{1}{2}})^{\frac{1}{2}}B^{\frac{1}{2}}.$$

It is known that $X_0 = B \sharp C$ is the unique positive solution of the Riccati equation $X^*B^{-1}X = C$, see [1,4,6,11].

Very recently, Izumino and Nakamura [5] gave some interesting consideration to weakly positive operators introduced by Wigner [12]. An operator T is weakly positive if $T = SCS^{-1}$ for some $S, C > 0$, where $X > 0$ means it is positive and invertible. It is equivalent to be of form $T = AB$ for some $A, B > 0$. (Take $A = S^2$ and $B = S^{-1}CS^{-1}$.) They pointed out that the square root $T^{\frac{1}{2}}$ of a weakly positive operator $T = SCS^{-1} = AB$ can be defined by $T^{\frac{1}{2}} = SC^{\frac{1}{2}}S^{-1} = A^{\frac{1}{2}}(A^{\frac{1}{2}}BA^{\frac{1}{2}})^{\frac{1}{2}}A^{-\frac{1}{2}}$, and

$$A^{-1} \sharp B = A^{-1}(AB)^{\frac{1}{2}}.$$

As an easy consequence, $A^{-1} \sharp B$ is a (unique) positive solution of Riccati equation $XAX = B$ for given $A, B > 0$.

Inspired by Ando's work (2) and Izumino-Nakamura's consideration, we would like to introduce a Riccati inequality by the positivity of an operator matrix, i.e.,

$$\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0 \quad \text{for given } B, A \geq 0$$

2. Riccati inequality.

In this section, we investigate solutions of Riccati inequality, in which we characterize them by factorization.

The following lemma is well-known, but important.

Lemma 1. *Let A be a positive operator. Then*

$$\begin{pmatrix} 1 & X \\ X^* & A \end{pmatrix} \geq 0 \quad \text{if and only if} \quad A \geq X^*X.$$

Proof. Since

$$\begin{pmatrix} 1 & 0 \\ 0 & A - X^*X \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -X^* & 1 \end{pmatrix} \begin{pmatrix} 1 & X \\ X^* & A \end{pmatrix} \begin{pmatrix} 1 & -X \\ 0 & 1 \end{pmatrix},$$

it follows that

$$\begin{pmatrix} 1 & X \\ X^* & A \end{pmatrix} \geq 0 \quad \text{if and only if} \quad A \geq X^*X.$$

We here note the existence of the maximum of the geometric mean (2), as an application of an idea in Lemma 1: We may assume that B is invertible. Then

$$\begin{pmatrix} B & X \\ X^* & C \end{pmatrix} \geq 0 \quad \text{if and only if} \quad C \geq X^*B^{-1}X$$

because

$$\begin{pmatrix} 1 & 0 \\ -X^*B^{-1} & 1 \end{pmatrix} \begin{pmatrix} B & X \\ X^* & C \end{pmatrix} \begin{pmatrix} 1 & -B^{-1}X \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} B & 0 \\ 0 & C - X^*B^{-1}X \end{pmatrix}.$$

Hence, if $C \geq X^*B^{-1}X$, then

$$B^{-\frac{1}{2}}CB^{-\frac{1}{2}} \geq (B^{-\frac{1}{2}}XB^{-\frac{1}{2}})^2.$$

By Lowner-Heinz inequality, we have

$$(B^{-\frac{1}{2}}CB^{-\frac{1}{2}})^{\frac{1}{2}} \geq B^{-\frac{1}{2}}XB^{-\frac{1}{2}},$$

so that

$$B \sharp C = B^{\frac{1}{2}}(B^{-\frac{1}{2}}CB^{-\frac{1}{2}})^{\frac{1}{2}}B^{\frac{1}{2}} \geq X.$$

Consequently, the maximum

$$\max\{X \geq 0; \begin{pmatrix} B & X \\ X^* & C \end{pmatrix} \geq 0\} = \max\{X \geq 0; C \geq X^*B^{-1}X\}$$

is given by $B \sharp C$.

Lemma 2. *Let A and B be positive operators. Then*

$$\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0 \quad \text{implies} \quad \text{ran} W \subseteq \text{ran} B^{\frac{1}{2}}.$$

and so $X = B^{-\frac{1}{2}}W$ is well-defined as a mapping.

Proof. Let $S = \begin{pmatrix} a & b \\ b^* & d \end{pmatrix}$ be the square root of $R = \begin{pmatrix} B & W \\ W^* & A \end{pmatrix}$. Then

$$R = S^2 = R = \begin{pmatrix} a^2 + bb^* & ab + bd \\ b^*a + db^* & b^*b + d^2 \end{pmatrix},$$

that is,

$$B = a^2 + bb^* \text{ and } W = ab + bd.$$

Therefore $\text{ran } B^{\frac{1}{2}}$ contains both $\text{ran } a$ and $\text{ran } b$, so that it contains $\text{ran } a + \text{ran } b$. Moreover $\text{ran } W$ is contained in $\text{ran } a + \text{ran } b$ by $W = ab + bd$.

Under the preparation of Lemmas 1 and 2, Riccati inequality can be solved as follows:

Theorem 3. Let A and B be positive operators on K and H respectively, and W be an operator from K to H . Then $\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0$ if and only if $W = B^{\frac{1}{2}}X$ for some operator X from K to H and $A \geq X^*X$.

Proof. Suppose that $\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0$. Since $\text{ran } W \subseteq \text{ran } B^{\frac{1}{2}}$ by Lemma 2, Douglas' majorization theorem [3] says that $W = B^{\frac{1}{2}}X$ for some operator X . Moreover we restrict X by $P_B X = X$, where P_B is the range projection of B . Noting that $y \in \text{ran } B$ if and only if $y = B^{\frac{1}{2}}x$ for some $x \in \text{ran } B^{\frac{1}{2}}$, the assumption implies that

$$\left(\begin{pmatrix} P_B & X \\ X^* & A \end{pmatrix} \begin{pmatrix} y \\ z \end{pmatrix}, \begin{pmatrix} y \\ z \end{pmatrix} \right) = \left(\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \begin{pmatrix} x \\ z \end{pmatrix}, \begin{pmatrix} x \\ z \end{pmatrix} \right) \geq 0$$

for all $y \in \text{ran } B$ and $z \in K$. This means that $\begin{pmatrix} P_B & X \\ X^* & A \end{pmatrix} \geq 0$, and so

$$\begin{pmatrix} P_B & 0 \\ 0 & A - X^*X \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ -X^* & A \end{pmatrix} \begin{pmatrix} P_B & X \\ X^* & A \end{pmatrix} \begin{pmatrix} 1 & -X \\ 0 & 1 \end{pmatrix} \geq 0,$$

that is, $A \geq X^*X$, as required. The converse is easily checked.

The following factorization theorem [2; Theorem I.1] is led by Theorem 3 and Douglas' factorization theorem [3].

Theorem 4. Let A and B be positive operators. Then $\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0$ if and only if $W = B^{\frac{1}{2}}VA^{\frac{1}{2}}$ for some contraction V .

Proof. Suppose that $\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0$. Then it follows from Theorem 3 that $W = B^{\frac{1}{2}}X$ for some bounded X satisfying $A \geq X^*X$. Hence we can find a contraction V with $X = VA^{\frac{1}{2}}$ by [3], so that $W = B^{\frac{1}{2}}VA^{\frac{1}{2}}$ is shown.

The converse is proved by Lemma 1 as follows:

$$\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} = \begin{pmatrix} B & B^{\frac{1}{2}}VA^{\frac{1}{2}} \\ A^{\frac{1}{2}}V^*B^{\frac{1}{2}} & A \end{pmatrix} = \begin{pmatrix} B^{\frac{1}{2}} & 0 \\ 0 & A^{\frac{1}{2}} \end{pmatrix} \begin{pmatrix} 1 & V \\ V^* & 1 \end{pmatrix} \begin{pmatrix} B^{\frac{1}{2}} & 0 \\ 0 & A^{\frac{1}{2}} \end{pmatrix} \geq 0.$$

3. An exact expression of the harmonic mean. Recall that the harmonic mean is defined by

$$B ! C = \max\{X \geq 0; \begin{pmatrix} 2B & 0 \\ 0 & 2C \end{pmatrix} \geq \begin{pmatrix} X & X \\ X & X \end{pmatrix}\}.$$

To obtain the exact expression of the harmonic mean, we need the following lemma which is more explicit than Theorem 3.

Lemma 5. *If $\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \geq 0$, then $X = B^{-\frac{1}{2}}W$ is bounded and $A \geq X^*X$.*

Proof. For a fixed vector x , we put $x_1 = B^{-\frac{1}{2}}Wx$. Since $B^{\frac{1}{2}}x_1 = Wx$, we may assume $x_1 \in (\ker B^{\frac{1}{2}})^\perp$. So it follows that

$$\begin{aligned} \|B^{-\frac{1}{2}}Wx\| &= \sup\{|(Wx, v)|; \|v\| = 1\} \\ &= \sup\{|(B^{-\frac{1}{2}}Wx, B^{\frac{1}{2}}u)|; \|B^{\frac{1}{2}}u\| = 1\} \\ &= \sup\{|(Wx, u)|; (Bu, u) = 1\}. \end{aligned}$$

On the other hand, since

$$\left(\begin{pmatrix} B & W \\ W^* & A \end{pmatrix} \begin{pmatrix} u \\ tx \end{pmatrix}, \begin{pmatrix} u \\ tx \end{pmatrix} \right) = |t|^2(Ax, x) + 2\operatorname{Re} t(Wx, u) + (Bu, u) \geq 0$$

for all scalars t , we have

$$|(Wx, u)|^2 \leq (Ax, x)(Bu, u).$$

Hence it follows that

$$\|B^{-\frac{1}{2}}Wx\|^2 = \sup\{|(Wx, u)|; (Bu, u) = 1\} \leq (Ax, x),$$

which implies that $X = B^{-\frac{1}{2}}W$ is bounded and $A \geq X^*X$.

Theorem 6. *Let B, C be positive operators. Then*

$$B ! C = 2(B - [(B + C)^{-\frac{1}{2}}B]^*[(B + C)^{-\frac{1}{2}}B]).$$

In particular, if $B + C$ is invertible, then

$$B ! C = 2(B - B(B + C)^{-1}B) = 2B(B + C)^{-1}C.$$

Proof. First of all, the inequality $\begin{pmatrix} 2B & 0 \\ 0 & 2C \end{pmatrix} \geq \begin{pmatrix} X & X \\ X & X \end{pmatrix}$ is equivalent to

$$\begin{pmatrix} 2(B + C) & -2B \\ -2B & 2B - X \end{pmatrix} = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2B - X & -X \\ -X & 2C - X \end{pmatrix} \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix} \geq 0.$$

Then it follows from Lemma 5 that $D = [2(B + C)]^{-\frac{1}{2}}(-2B)$ is bounded and $D^*D \leq 2B - X$. Therefore we have the explicit expression of $B ! C$ even if both B and C are non-invertible:

$$B ! C = \max\{X \geq 0; D^*D \leq 2B - X\} = 2B - D^*D.$$

In particular, if $B + C$ is invertible, then

$$B ! C = 2B - D^*D = 2(B - B(B + C)^{-1}B) = 2B(B + C)^{-1}C.$$

4. Pedersen-Takesaki theorem. Finally we review a work of Pedersen and Takesaki [8] from the viewpoint of Riccati inequality; we add another equivalent condition to their theorem:

Theorem 7. *Let B and C be positive operators and B be nonsingular. Then the following statements are mutually equivalent:*

- (1) *Riccati equation $XBX = C$ has a positive solution.*
- (2) *$(B^{\frac{1}{2}}CB^{\frac{1}{2}})^{\frac{1}{2}} \leq kB$ for some $k > 0$.*
- (3) *There exists the minimum of $\{X \geq 0; C \leq XBX\}$.*
- (3') *There exists the minimum of $\{X \geq 0; \begin{pmatrix} 1 & C^{\frac{1}{2}} \\ C^{\frac{1}{2}} & XBX \end{pmatrix} \geq 0\}$.*

Proof. We first note that (3) and (3') are equivalent by Lemma 1.

Now we suppose (1), i.e., $X_0BX_0 = C$ for some $X_0 \geq 0$. If $X \geq 0$ satisfies $C \leq XBX$, then

$$(B^{\frac{1}{2}}X_0B^{\frac{1}{2}})^2 = B^{\frac{1}{2}}CB^{\frac{1}{2}} \leq (B^{\frac{1}{2}}XB^{\frac{1}{2}})^2$$

and so

$$B^{\frac{1}{2}}X_0B^{\frac{1}{2}} \leq B^{\frac{1}{2}}XB^{\frac{1}{2}}.$$

Since B is nonsingular, we have $X_0 \leq X$, namely (3) is proved.

Next we suppose (3). Since $C \leq XBX$ for some X , we have

$$B^{\frac{1}{2}}CB^{\frac{1}{2}} \leq (B^{\frac{1}{2}}XB^{\frac{1}{2}})^2$$

and so

$$(B^{\frac{1}{2}}CB^{\frac{1}{2}})^{\frac{1}{2}} \leq B^{\frac{1}{2}}XB^{\frac{1}{2}} \leq \|X\|B,$$

which shows (2).

The implication (2) \rightarrow (1) has been shown by Pedersen-Takesaki [8] and Nakamoto [7], but we sketch it for convenience. By Douglas' majorization theorem [3], we have

$$(B^{\frac{1}{2}}CB^{\frac{1}{2}})^{\frac{1}{2}} = ZB^{\frac{1}{2}}$$

for some Z , so that

$$(B^{\frac{1}{2}}CB^{\frac{1}{2}})^{\frac{1}{2}} = B^{\frac{1}{2}}Z^*ZB^{\frac{1}{2}} \quad \text{and} \quad B^{\frac{1}{2}}CB^{\frac{1}{2}} = B^{\frac{1}{2}}(Z^*ZBZ^*Z)B^{\frac{1}{2}}.$$

Since B is nonsingular, Z^*Z is a solution of $XBX = C$.

Acknowledgement. We remark that similar discussion to ours in Section 2 has been done by several authors, e.g. [9] and [10]. The authors would like to express their thanks to Professor Tomiyama and Professor Kosaki for thier kind suggestions on the above.

REFERENCES

- [1] W.N.ANDERSON, JR. AND G.E.TRAPP, *Operator means and electrical networks*, Proc. 1980 IEEE International Symposium on Circuits and systems, 1980, 523-527.
- [2] T.AND0, *Topics on Operator Inequalities*, Lecture Note, Sapporo, 1978.
- [3] R.G.DOUGLAS, *On majorization, factorization, and range inclusion of operators on Hilbert space*, Proc. Amer. Math. Soc., 17(1966), 413-415.
- [4] J.I.FUJII AND M.FUJII, *Some remarks on operator means*, Math. Japon., 24(1979), 335-339.
- [5] S.IZUMINO AND M.NAKAMURA, *Wigner's weakly positive operators*, Sci. Math. Japon., to appear.
- [6] F.KUBO AND T.AND0, *Means of positive linear operators*, Math. Ann., 246(1980), 205-224.
- [7] R.NAKAMOTO, *On the operator equation $THT = K$* , Math. Japon., 24(1973), 251-252.
- [8] G.K.PEDERSEN AND M.TAKESAKI, *The operator equation $THT = K$* , Proc. Amer. Math. Soc., 36(1972), 311-312.
- [9] L.M.SCHMITT, *The Radon-Nikodym theorem for L^p -spaces of W^* -algebras*, Publ. RIMS, Kyoto Univ., 22(1986), 1025-1034.
- [10] JU.L.ŠMUL'JAN *An operator Hellinger integral*, Mat. Sb., 91(1959), 381-430. (English translation: *A Hellinger operator integral*, Amer. Math. Soc. Translations, Ser. 2, 22(1962), 289-338.)
- [11] G.E.TRAPP, *Hermitian semidefinite matrix means and related matrix inequalities - An introduction*, Linear Multilinear Alg., 16(1984), 113-123.
- [12] E.P.WIGNER, *On weakly positive operators*, Canadian J. Math., 15(1965), 313-317.

Department of Information Sciences, Osaka Kyoiku University, Asahigaoka, Kashiwara, Osaka 582-8582, Japan.

fujii@cc.osaka-kyoiku.ac.jp

Department of Mathematics, Osaka Kyoiku University, Asahigaoka, Kashiwara, Osaka 582-8582, Japan.

mfujii@cc.osaka-kyoiku.ac.jp

Faculty of Engineering, Ibaraki University, Hitachi, Ibaraki 316-8511, Japan.

nakamoto@base.ibaraki.ac.jp